

A&A manuscript no.
(will be inserted by hand later)

Your thesaurus codes are:
02.09.1, 02.13.1, 13.25.5, 08.02.3, 08.23.3, 08.09.2: AM Herculis

ASTRONOMY
AND
ASTROPHYSICS

Magnetic field and unstable accretion during AM Herculis low states. ^{*}

J.M. Bonnet-Bidaud¹, M. Mouchet^{2,3}, N.M. Shakhovskoy⁴, T.A. Somova⁵, N.N. Somov⁵, I.L. Andronov⁶, D. de Martino⁷, S.V. Kolesnikov⁶, and Z. Kraicheva⁸

¹ Service d'Astrophysique, DSM/DAPNIA/SAP, CE Saclay, F-91191 Gif sur Yvette Cedex, France

² DAEC, Observatoire de Paris Section de Meudon, F-92195 Meudon Cedex, France

³ Université Denis Diderot, Place Jussieu, F-75005 Paris, France

⁴ Crimean Astrophysical Observatory, Nauchnyi, Crimea, 334413, Ukraine

⁵ Special Astrophysical Observatory, Nizhnij Arkhyz, Karachaj-Cherkess Republik, 357147, Russia

⁶ Department of Astronomy, Odessa St University, Shevchenko Park, Odessa, 270014, Ukraina

⁷ Osservatorio Astronomico di Capodimonte, I-80131 Napoli, Italy

⁸ Institute of Astronomy, Bulgarian Academy of Sciences, 72 Tsarigradsko Shouse Blvd., 1784 Sofia, Bulgaria

Received date: 29 September 1999; accepted date: 23 December 1999

Abstract. A study of AM Her low states in September 1990 and 1991 and June-July 1997 is reported from a coordinated campaign with observations obtained at the Haute-Provence observatory, at the 6-m telescope of the Special Astrophysical Observatory and at the 2.6m and 1.25m telescopes of the Crimean observatory. Spectra obtained at different dates when the source was in low states at a comparable V magnitude, show the presence of strong Zeeman absorption features and marked changes in emission lines with a day-to-day reappearance of the HeII (4686Å) emission lines in 1991. Despite this variability, the magnetic field inferred from the fitting of the absorption spectrum with Zeeman hydrogen splitting, is remarkably constant with a best value of (12.5 ± 0.5) MG. Detailed analysis of the UBVRI light curves shows the presence of repetitive moderate amplitude (~ 0.3 - 0.5 mag) flares predominantly red in colour. These flares are attributed to small accretion events and are compared to the large (~ 2 mag.) blue flare reported by Shakhovskoy et al. (1993). We suggest that the general flaring activity observed during the low states is generated by accretion events. The different characteristics of the flares (colour and polarization) are the results of different shock geometries depending on the net mass accretion flux.

Key words: stars: white dwarf - stars: accretion - AM Herculis - magnetic field

1. Introduction

AM Herculis (4U1814+49) is the well-known prototype of the "polar" systems, a subclass of cataclysmic variables in which a highly magnetized (10^7 G) white dwarf in a close binary system accretes matter from a low-mass companion (see Cropper 1990, Chanmugam 1992 for a review). From the long term optical monitoring collected and kindly made available to us by the AAVSO (J. Mattei, private communication), it is now well known to oscillate irregularly between two different states of optical brightness, a high state at ($V \sim 12.5$) corresponding to high accretion rate and a low state ($V \sim 15$) during which the accretion luminosity is significantly reduced so that photospheric emission of the two stars becomes visible in the infrared for the companion and in the UV and the optical for the white dwarf. The low state reveals in particular a complicated optical spectrum where strong absorption features due to the Zeeman splitting of the Balmer lines produced in the high surface magnetic field of the white dwarf are clearly visible, allowing the direct measurement of the field (Schmidt et al. 1981, Latham et al. 1981, Young et al. 1981). During low states, the UV emission is also found consistent with white dwarf atmosphere models with $T = (2-2.5) \cdot 10^4$ K and typical size $(6-8) \cdot 10^6$ m (Heise & Verbunt 1988, Gnsicke et al. 1995, Silber et al. 1996).

The question of whether or not the accretion ceases during low states is still open. UV data indicate that the polar caps are still substantially heated and the few low states observed in the X-rays reveal the presence of residual accretion. A weak X-ray modulation due to the occultation of the main accreting pole is detected (Fabbiano 1982, de Martino et al. 1998). At such low rates, the accretion onto the white dwarf is highly unstable and eventually switches-off as recently observed by the BeppoSAX satel-

Send offprint requests to: J.M. Bonnet-Bidaud

^{*} Based on data collected at SAO (Russia), CAO (Crimea), BAO (Bulgaria) and OHP (France)

Correspondence to: bobi@discovery.saclay cea.fr

lite (de Martino et al. 1998). Strong (30%) quasi-periodic optical oscillations with periods near 5 minutes have been observed during a decline to a low state and interpreted as an accretion instability arising close to the capture radius (Bonnet-Bidaud et al. 1991). A spectacular sharp rising (~ 1 hr) flare of ~ 2 mag. was also detected during a 1992 low state, tentatively associated with a stellar flare from the red dwarf companion (Shakhovskoy et al. 1993). We present here data obtained during different low states of AM Her in 1990, 1991 and 1997 which show that the source presents very different characteristics and variability. The 1997 data were obtained to supplement contemporaneous BeppoSAX X-ray observations (de Martino et al. 1998).

2. Observations

2.1. Photometry and polarimetry

Photometric and polarimetric observations were conducted at the 1.25m AZT-11 telescope of the Crimean Astrophysical Observatory (CAO), equipped with the double-beam chopping polarimeter of the Helsinki University (Korhonen et al. 1984). On 1991 Sept 4 and 5 and 1997 July 1 and 30, UBVRI data were collected during ~ 3 -4 h intervals overlapping spectroscopic observations described below. The UBVRI photometric data were recorded automatically with a 23.1s resolution and the polarimeter was used in circular polarization mode with a resolution of 3 min in the same bands. However a statistically significant signal was only received in circular polarization in the R and I bands and only the corresponding data are discussed here. The full description of the polarimeter and the method of the observations are presented in Berdyugin & Shakhovskoy (1993). On 1997 July 1, data were also obtained at the CAO 2.6m Shajn telescope, using a one-channel polarimeter with a fast rotating achromatic quarter-wave plate, in the wide R-band (0.50-0.75nm) with a 4s integration time.

On 1991 Sept 4 and 5, photometric data were acquired at the 6-m BTA telescope of the Special Astrophysical Observatory (SAO) (Nizhnij Arkhyz, Russia). The observations were carried out at the Nasmyth focus of the telescope, simultaneously with the spectroscopy, using 50% of the incoming flux split by a dichroic plate to the NEF photometer (Vikuliev et al. 1991). A light curve through a Johnson B-filter with a 12 arcsec aperture was recorded with a 0.1s resolution during the observations. UBVR measurements were also performed at the beginning and end of the observations to calibrate the brightness level of the source. On 1997 July 3 and 4, complementary photometric data were also obtained at the Belgradchik Astronomical Observatory BAO (Bulgaria), using an ST-8 CCD camera attached to a 60cm telescope.

All UBVR magnitudes were obtained from differential measurements, using the star D ($m_V = 13.1$) in the field

as a comparison (Liller 1977, Andronov & Korotin, 1982).

2.2. Spectroscopy

Spectroscopic data were collected during AM Her low states in 1990, 1991 and 1997 using the SP-124 spectrograph of the 6-m BTA telescope (Ioannisianni et al. 1982). A television scanner with two lines of 1024 channels is used to record the sky and source spectra simultaneously in a photon-counting mode (see Somova et al. 1982, for a detailed description of the instrumentation). A 2.5 arcsec aperture was set, adapted to the seeing and the spectrograph was equipped with a grating yielding a resolution of $4\text{\AA}$, $2\text{\AA}$ and $2\text{\AA}$, respectively in 1990, 1991 and 1997. Data were reduced using the SIPRAN software developed at SAO (Somov 1986). Part of the 1990 spectroscopic data was already preliminarily discussed in Bonnet-Bidaud et al. (1992).

On 1997, spectroscopic observations were also obtained at the 1.93m telescope of the Haute-Provence Observatory (OHP), equipped with the Carelec spectrograph (Lemaitre et al. 1990).

The log of the observations is presented in Table 1 with the Heliocentric Julian Dates (HJD) corresponding to the start of the exposures. In the following, the orbital/rotational phases have been computed according to the ephemeris derived by S. Tapia and quoted in Heise & Verbunt (1988), where $\text{HJD}(\phi = 0) = 2443014.76614(4) + 0.128927041(5)\text{E}$, $\phi = 0$ corresponding to the maximum of linear polarization.

3. Analysis and results

3.1. Photometry and polarimetry

Figures 1 and 2 show the UBVRI light curves covering more than one orbital cycle in the different observations. The source is seen at a mean V magnitude of ~ 15.1 and ~ 15.2 , respectively in 1991 and 1997, consistent with the level previously reported during low states of the star (Szkody et al. 1982, Bailey et al. 1988). Table 2 gives the mean magnitude and dispersion in the different bands and at the different epochs.

In 1991, the shape of the modulation is typical of a low state with a broad hump in U and B around phase 0.6, antiphased with a minimum in R and I. A secondary minimum in R and I is also visible around phase 0.2. The V light curve is flat but with a greater dispersion. Superimposed on the smooth modulation, clear rapid (5-10 min.) flux enhancements are also seen, particularly visible in R and I bands. Characteristics of these flares (marked with numbers on Figure 1) are analysed below. In 1997, the source is less active, though at a similar brightness level with no evidence of flares. For all observing nights, the modulation is nearly absent in all bands and no strong

Table 1. Journal of observations.

<i>Date</i>	<i>Telescope</i>	<i>Observations</i>	<i>Range</i>	<i>Resolution</i>	<i>HJD start</i>	<i>Exp(min)</i>
1990 Sep. 17	SAO 6m	spectroscopy	3800-5800Å	4Å	8152.2342	144
1991 Sep. 04	SAO 6m	spectroscopy	3950-4950Å	2Å	8504.2340	102
"	SAO 6m	photometry	B	0.1s	8504.2405	83
"	CAO 1.25m	photom./polarim.	UBVRI	23.1s/180s	8504.2497	224
1991 Sep. 05	SAO 6m	spectroscopy	3950-4950Å	2Å	8505.2330	85
"	SAO 6m	photometry	B	0.1s	8505.2488	61
"	CAO 1.25m	photom./polarim.	UBVRI	23.1s/180s	8505.2543	243
1997 June. 30	SAO 6m	spectroscopy	4050-5070Å	2Å	10630.4583	196
1997 July 01	OHP 1.93m	spectroscopy	3750-7250Å	13Å	10631.3647	90
	CAO 2.6m	photom./polarim.	R	4s/128s	10631.3329	244
	CAO 1.25m	photom./polarim.	UBVRI	20s/192s	10631.3711	182
1997 July 03	BAO 0.6m	photometry	V	60s	10633.4066	225
1997 July 04	BAO 0.6m	photometry	V	60s	10634.4074	242
1997 July 29	OHP 1.93m	spectroscopy	3750-7250Å	13Å	10659.3967	180
1997 July 30	CAO 1.25m	photom./polarim.	UBVRI	23.1s/180s	10660.3166	177
1997 July 31	OHP 1.93m	spectroscopy	3750-7250Å	13Å	10660.5985	45
"	CAO 1.25m	photom./polarim.	UBVRI	23.1s/180s	10661.3513	82

Table 2. Magnitudes in quiescence.

<i>Date</i>		<i>U</i>		<i>B</i>		<i>V</i>		<i>R</i>		<i>I</i>
1992 Aug 29(*)	Q92	14.90		15.58		15.40		14.79		13.56
1991 Sep. 4	Q4	14.72	(0.11)	15.52	(0.09)	15.11	(0.17)	14.49	(0.19)	13.41 (0.18)
1991 Sep 5	Q5	14.57	(0.13)	15.38	(0.08)	15.08	(0.20)	14.31	(0.21)	13.32 (0.20)
1997 July 1	Q1	14.82	(0.12)	15.55	(0.11)	15.34	(0.24)	14.70	(0.08)	13.60 (0.11)
1997 July 30	Q30	14.81	(0.12)	15.53	(0.09)	15.16	(0.22)	14.61	(0.19)	13.44 (0.22)

(*) from Shakhovskoy et al. (1993)

Numbers in parentheses are a measure of the dispersion

flux variations are seen, except for a dip around phase 0.1 in I on July 1 and a ~ 0.3 mag. broad increase in the R-band near phase 0. on July 30. BAO light curves obtained on July 3 and 4 also show no modulation.

Figure 3a and b shows the corresponding circular polarization in the R and I bands. A significant polarization is seen in 1991, with a mean level in the R / I bands of -3.9% / -2.6% and -4.5% / -4.0% on Sept 4 and 5 respectively, indicating a significant residual accretion. Negative polarization reaches a maximum of up to -12% around phase 0.4 and 0.7 and a minimum near 0% around phase 0.1, comparable to what usually observed during normal high states of the source (Bailey & Axon 1981). The polarization, attributed to cyclotron radiation, is usually restricted to the infrared bands during low state observations and shifted to higher frequencies only during high states (Bailey et al. 1988). It is usually absent in R and I bands during low states (Shakhovskoy et al. 1993, Silber et al. 1996) though occasional detections have been made (Latham et al. 1981, Shakhovskoy et al. 1992). In 1997, at a comparable brightness level, no significant polarization is detected in the same R and I bands, with mean values respectively of $(0.8 \pm 1.9)\%$ and $(0.7 \pm 1.8)\%$ for July 1 and $(1.9 \pm 4.2)\%$

and $(0.5 \pm 2.3)\%$ for July 30, indicating that the cyclotron emission has become negligible (Shakhovskoy et al. 1992).

The peculiar flaring variability of the source seen in 1991 has been investigated by computing the characteristics of the flares marked in Figure 1. Figure 4a shows the colour-colour diagram of the AM Her system during the quiescent state (outside flares) and at the peak of the flares (with the size of symbols indicative of the flare intensity). For comparison, the colours of the very large blue flare observed by Shakhovskoy et al. (1993) during a 1992 low state are also shown. The AM Her colours during quiescence are remarkably similar in the different epochs. The flares are clearly distributed into three different categories: moderate flares (F3,F5,F6) during which the system stays approximately of the same colour as in the quiescent state or slightly bluer, intense flares (F1,F2,F4) during which the colours change to strongly red and the very intense 1992 flare clearly peaking in the blue.

The energy distribution of the flares has been evaluated by computing in the different bands the difference between the magnitude at peak and the local quiescent magnitude, estimated from polynomial fits through the light curve excluding flares. Magnitude excess due to the flares is reported in Table 3 and the corresponding spec-




Fig. 1. UBVRI light curves of the AM Her low state in 1991. (left Sept. 4, right Sept. 5). Clear flaring activity is seen. The strongest flares have been marked by numbers

tra of selected flares are given in Figure 4b. The slope of the best linear fit to the $(\log F_\nu - \log \nu)$ distribution is also given in Table 3. As indicated by the slope, the strongest flares appear clearly red, peaking around R/I bands with a slope $\sim \nu^{-(2-3)}$, while the less intense ones are bluer $\sim \nu^{-(1)}$. We note that the polarization during the largest flares is significant in both R and I bands (see open symbols in Fig. 3a and b). The properties of the flares are further discussed in Section 4.

Quasi-periodic oscillations (QPO) have been searched using (0.1s) resolution photometric data obtained in 1991. While no significant QPOs have been found on 1991 September 4, QPOs are clearly detected on 1991 September 5, with an amplitude 8.6% and a period 6.6 min (397s) during 30 minutes covering a (0.97-1.14) phase interval. No QPOs were observed in 1997.

3.2. Spectroscopy

The spectra obtained during the 1990, 1991 and 1997 low states have been averaged to produce a mean normalized spectrum representative of each different epoch (Figure 5). The spectra were reduced by standard procedure using MIDAS-ESO package and fluxes have been normalized by dividing by a continuum fitted through selected points free of line features. All spectra show clear evidence of the Balmer lines in emission with their associated Zeeman components in absorption, similar to what previously observed (Latham et al. 1981, Schmidt et al. 1981, Young et al. 1981, Silber et al. 1996). The equivalent widths of the emission lines appear to vary in the different spectra (see Table 4). The HeII 4686 line is only detectable in 1991. At this epoch, the spectrum changes drastically from one day to the next, with a sudden appearance of this high excitation line, and an equivalent width changing from $(0.4 \pm 0.5)\text{\AA}$ to $(3.6 \pm 0.5)\text{\AA}$. On 1991 Sept. 5, though the system is at the same mean low level ($m_V = 15.1$) and




Fig. 2. UBVR light curves of the AM Her low state in 1997 (left July 1, right July 30). No clear modulation is seen except for the flux increase at the end of July 30

Table 3. Magnitude excess due the flares with respect to the local quiescence level.

<i>Time(HJD)</i>	<i>Flare</i>	<i>U</i>	<i>B</i>	<i>V</i>	<i>R</i>	<i>I</i>	$\alpha(F_{\nu} \sim \nu^{\alpha})$
8864.3612	F92	-2.54	-2.36	-2.14	-1.71	-0.84	+0.82 (0.14)
8504.3862	F1	-0.32	-0.37	-0.67	-0.77	-0.63	-2.75 (0.26)
8505.2875	F2	-0.13	-0.13	-0.24	-0.31	-0.30	-3.09 (0.30)
8505.3400	F3	-0.37	-0.35	-0.38	-0.33	-0.20	-0.53 (0.17)
8505.3799	F4	-0.15	-0.13	-0.43	-0.47	-0.35	-3.29 (0.59)
8505.3903	F5	-0.26	-0.28	-0.53	-0.23	-0.24	-1.19 (0.45)
8505.4135	F6	-0.20	-0.25	-0.27	-0.44	-0.42	-2.79 (0.36)

Error bars are given in parentheses

the Zeeman absorption lines, typical of low state, are still clearly present, the emission lines are strong with H_{β} and H_{γ} EW of $19\text{\AA}$ and $14\text{\AA}$ respectively, within a factor 1.5-2 of the high state values. This suggests a significant residual accretion.

The location and depth of the Zeeman absorption features are best shown in Figure 6, where the mean of all low and medium resolution spectra are displayed on an extended

scale, with the emission part being cut. The two sets of data appear very similar with, at higher resolution, the evidence of clear and rather sharply defined absorptions around 4080, 4300, 4650 and $4820\text{\AA}$. To identify these features, synthetic idealized spectra have been constructed using the Zeeman wavelengths and oscillator strengths tabulated by Kemic (1974) for different magnetic fields in the range of 10-30MG as expected for AM Her. The in-




Fig. 3. UBVR circular polarization light curves in AM Her low states (left July 04 (top) and 05 (bottom), right July 01 (top) and 30 (bottom)). Open symbols in 1991 correspond to the flares marked in Figure 1

tensities of the lines were taken to be proportional to the oscillator strengths in the Milne-Eddington approximation (see Latter et al. 1987) and the spectra were interpolated with respect to the field strength to provide a complete grid of comparison spectra. The computed "theoretical" spectra were further convolved with an instrumental response corresponding to a spectral resolution of $2\text{\AA}$.

The inspection of the synthetic spectra reveals that the most significant features of Fig. 6 correspond to the σ and π components of the H_β and H_γ hydrogen lines. In the range of considered B fields, these lines clearly distributed into "stationary" lines, only weakly variable in position, such as $H_\beta \sigma^+$ and π and "non stationary lines" strongly or moderately variable in position such as $H_\beta \sigma^-$ and $H_\gamma \pi$ (Angel 1978). The non stationary lines allow a precise determination of the magnetic field strength. In Fig. 6 is shown the $B=12.5\text{MG}$ synthetic spectrum which is the best description of the data. Most features are accurately reproduced such as the split components of H_β and H_γ . The accuracy in the B determination is typically $\pm 0.5\text{MG}$

based on a precision better than $\sim 10\text{\AA}$ in the position of the fast varying $H_\beta \sigma^-$ feature.

More careful inspection reveals two additional features at $\sim 4400\text{\AA}$ and $\sim 4775\text{\AA}$ that are not reproduced by the synthetic spectrum. Interestingly enough, the $\sim 4775\text{\AA}$ feature, which is clearly visible in Fig. 6 as a left shoulder to the $H_\beta \pi$ feature, can be reproduced by a $H_\beta \pi$ component from a significant higher field ($\sim 17\text{MG}$). The corresponding $H_\beta \sigma^-$ will then be shifted to $\sim 4560\text{\AA}$ where a small feature is indeed present in the spectrum so we cannot exclude the possible superposition of this higher field. A similar additional higher field component has already been reported by Latham et al. (1981). We also investigated the possible contributions from helium lines. For fields in the polar range, Zeeman helium stationary components are expected from the HeI $4471\text{\AA}$ line at 4320 , 4420 and $4530\text{\AA}$, not in accordance with the observed features.

We looked for possible variations of the main Zeeman features with orbital phase using our longest set of data of




Fig. 4. a (top)- $(V-R/B-V)$ colour-colour diagram of AM Her during the flares (F) and during the quiescent (Q) low states. The size of the flare symbols are indicative of their intensities. b(bottom)- Energy distribution ($\log F_\nu - \log \nu$) of the light excess due to the flares.

Table 4. Emission line equivalent widths (in Å).

Date	H_α	H_β	H_γ	$HeII(4686\text{\AA})$
90 Sep 17	-	6.0(5)	2.0(5)	-
91 Sep 04	-	8.7(5)	4.8(6)	0.4 (5)
91 Sep 05	-	19.0(8)	14.1(3)	3.6 (5)
97 Jun 30	-	6.0(6)	1.4(3)	-
97 Jul 01	32(1)	4.1(5)	1.8(2)	-
97 Jul 29	39(1)	5.6(5)	2.8(3)	-
97 Jul 30	25(1)	2.5(6)	1.2(3)	-

Error bars on last digit are given in parentheses

1997 July 29 at low resolution. The spectrum appears remarkably stable in phase with only possible minor variations in intensities around the $H_\beta \sigma^-$ line.

4. Discussion




Fig. 5. Mean low state spectra for the different epochs. Note the strong increase in the emission lines from 1991 Sept. 4 to 5 and Zeeman absorptions at Balmer lines visible in all spectra.

4.1. The AM Her magnetic field

The best magnetic field derived from the Zeeman absorption components in the AM Her optical spectrum during different episodes of low state in 1990, 1991 and 1997 is $B = (12.5 \pm 0.5) \text{ MG}$, in accordance with what previously reported but with better accuracy (Latham et al. 1981, Schmidt et al. 1981, Young et al. 1981, Silber et al. 1996). The Zeeman split absorption components in polar low states are usually assumed to originate in the hot photosphere of the magnetic white dwarf whose contribution becomes dominant when the accretion ceases.

The relative good accuracy in the magnetic field determination is achieved by the stability of "non stationary" components such as $H_\beta \sigma^-$ which are very sensitive to the field strength. Despite the spectra are averaged over different orbital phase intervals and different epochs, the presence of such a stable feature is an indication of a rather homogeneous field. This situation is surprising since, in a simple dipole model, a range of 2 is expected between

the polar and equatorial field and a spread of B-values is therefore expected for an hemisphere seen at a given inclination (Saffer et al. 1989). The observed restricted range would therefore imply a nearly equator-on view with an inclination close to 90° and an observed B field peaking at $B_{\text{polar}}/2$.

4.2. Unstable accretion during the low states

The AM Her optical low states reported in this paper, though all showing a similar optical brightness with $m_v=(15.1-15.4)$, display different overall characteristics. Of particular interest is the behaviour of the source in 1991 when both a significant polarization and flaring activity is seen contrary to 1990 and 1997. Within the brightness history of the source (Mattei J., AAVSO, private communication), it may be significant that the 1991 observation is located toward the end of a rather slow decline from high to low state, while the 1990 and 1997 observations are both included in a prolonged long stable low state, lasting already for 5 and 2 months respectively.

The major flares observed in 1991 are predominantly red in nature when compared to quiescent state. This and the presence of a significant polarization point toward residual unstable accretion at that time. The presence of a low level ($\sim 5\text{min.}$) temporary QPOs usually observed during intermediate high states (Bonnet-Bidaud et al. 1992) and the reappearance of high excitation lines on Sept 5, 1991 further strengthen this conclusion.

The comparison of this flaring activity with the major flare observed in 1992, also during a low state, is interesting. It has been proposed that the sharp 1992 event is due to a stellar flare originating from a magnetic active secondary (Shakhovskoy et al. 1993). The shape as well as the colour changes of the flare were found consistent with what observed during strong flares from red dwarfs (Beskrovnaya et al. 1996). The total energy in the flare, though at the very extreme upper end of what usually observed in nearby UV Ceti systems, is comparable to the more energetic flares detected in open clusters (Shakhovskaya 1989). However flares of this type are also fairly repetitive while it has been observed only once in AM Her although the source has been intensively monitored. We note however that a flare at this amplitude may be lost during high states. We find significant that the maximum magnitude of the flare ($m_v \sim 12.5$) is of the order of the high state level so that this episod may also be interpreted alternatively as an unstable accretion event. The very peculiar colours of this flare (see Fig. 4a) may result in this case from inhomogeneous accretion with blobs buried into the white dwarf atmosphere if the density is high enough.

Following Frank et al. (1988), we estimate the critical density for buried shocks by equating the accreted gas ram pressure with the atmosphere pressure at an optical depth sufficient for efficient reprocessing, giving

$$\rho_{cr} = 1.42 M_{wd}^2 [R_{wd}^3 T_{wd} \ln(97 R_{wd} \rho_{cr})]^{-1}$$

where ρ_{cr} is the critical density in units of $10^{-6} \text{ g.cm}^{-3}$ and M_{wd} , R_{wd} and T_{wd} the mass, radius and temperature of the white dwarf in units of $M_\odot$, 10^9 cm and 10^5 K respectively. Assuming $T_{wd} \sim 0.2$ and $M_{wd} = 0.6$ (see Gnsicke et al. 1995) with a corresponding $R_{wd} = 0.9$ (Nauenberg 1972) gives a critical density of $\rho_{cr} = 0.85 \cdot 10^{-6} \text{ g.cm}^{-3}$. This has to be compared with the mean accreted density during steady high states $\rho_{high} = \dot{M}/(v_{ff}A)$ with a typical accretion rate $\dot{M} \sim 10^{16} \text{ g.s}^{-1}$, v_{ff} the free-fall velocity $\sim 10^8 \text{ cm.s}^{-1}$ and A the accreting area $\sim 10^{16} \text{ cm}^2$ which yields $\rho_{high} = 10^{-8} \text{ g.cm}^{-3}$.

To achieve blobby accretion, the density in the flare has then to be ~ 10 -100 times the steady accretion. This may be easily achieved if, for instance, the accretion during the temporary event takes place onto a (3-10) reduced accretion spot radius due to particular unstable capture conditions by the magnetic field at that time. The emerging radiation from such a blobby accretion has not yet been computed accurately but the radiation is expected to be thermalized inside the atmosphere and will be radiated at temperature closer to the white dwarf blackbody (Kuijpers & Pringle 1982). We then expect the optical spectrum of the flare to follow roughly a ($\sim \nu^{+(1-2)}$) dependency, in accordance with the observed colours. Such flares should be mostly visible in soft X-rays.

On the opposite, the smaller flares observed during low states may correspond to small accretion events during which the accretion rate only temporarily increased, leading to transient presence of a standard shock above the white dwarf and an associated cyclotron emission. Such small scale flares, as those observed in 1991, show indeed polarization and red colours ($\sim \nu^{-(2-3)}$) expected from the optically thin part of the cyclotron. During the typical low states, outside flares, the low density accretion flow may lead to the absence of a shock and heat the upper atmosphere by Coulomb collisions leading to the so called bombardment solution (Woelk & Beuermann 1992). In this last case, the bluer colors of the optical radiation during quiescence can be easily explained by the decrease (1991) or negligible (1997) cyclotron emission. This picture derived from the optical variability study of AM Her is in accordance with the conclusions drawn from the low state X-ray characteristics of the source (Ramsay et al. 1995). The reason of the unstable accretion during low states, leading to both large accretion event and/or small accretion instability, is still unresolved as well as the exact mechanism responsible for the low states.

5. Conclusion

The study of the temporal and spectral characteristics of AM Her during low states at different periods allowed different conclusions to be drawn.

The Zeeman spectral features shown by the source are surprisingly very stable although obtained through different parts of the orbital cycle and therefore with




Fig. 6. Low state spectra showing clear Zeeman absorptions. The normalized continuum has been cut to suppress the emission lines. The low resolution spectrum (top) corresponds to the 1997 low state only while the higher resolution one (middle) is an average of 1990, 1991 and 1997 low states. The main Zeeman identifications are shown. The fitted best ($B=12.5$ MG) synthetic spectrum is displayed at bottom and reproduces most of the strongest features. The arrows indicate unidentified features (see text)

different orientations with respect to the suspected dipole magnetic field. The magnetic field strength derived from the position of "non stationary" lines is (12.5 ± 0.5) MG which could represent an averaged field seen over the white dwarf surface. Additional features are seen which may originate from an higher field region.

The temporal optical variability of AM Her during low states is very rich, displaying occasional large blue flare events as well as repetitive smaller amplitude red flares. It is shown that the characteristics of all these flares can be explained by accretion events of different amplitudes. The large and unique event observed in 1992, though consistent with red dwarf flares, can also be tentatively explained by a large increase of the accretion rate coupled to a reduced accretion area which can lead to a sufficient increase in the density to produce a buried shock unstable accretion. The more frequent smaller amplitude flares are interpreted instead as small variable increases of the accretion rate.

References

- Andronov I., Korotin S.A., 1982, Astr. circular N 1223, 1.
- Angel J.R.P, 1978, ARA&A 16, 487
- Bailey J., Axon D.J., 1981, MN 194, 187
- Bailey J., Hough J.H., Wickramasinghe D.T., 1988, MN 233, 395
- Berdyugin A.V., Shakhovskoy N.M., 1993, Bull. Crimean Astrophys. Obs. 87, 122-134
- Beskrovnaya N.G., Ikhsanov N.R., Bruch A., Shakhovskoy N.M., 1996, A&A 307, 840
- Bonnet-Bidaud J.M., Somova T., Somov N., 1991, A&A 251, L27
- Bonnet-Bidaud J.M., Somov N.N., Somova T.A., 1992, in Stellar magnetism, Eds Glagolevskij YU.V., Romanyuk I.I., p.181.
- Chanmugam G., 1992, ARA&A 30, 143
- Cropper M., 1990, Space Sci. Rev. 54, 195
- de Martino D., Gnsicke B.T., Matt G. et al., 1998, A&A 333, L31
- Fabbiano G., 1982, ApJ 262, 709
- Frank J., King A.R., Lasota J.P., 1988, A&A 193, 113
- Gnsicke B.T., Beuermann K., de Martino D., 1995, A&A 303, 127

- Heise J., Verbunt F., 1988, A&A, 189, 112
- Ioannisianni B.K. et al., 1982, in Instrumentation for astronomy with Large Optical Telescopes, Ed. C.H.Hamphries, Reidel, p.3
- Kemic S.B., 1974, in Joint Institute for Laboratory Astrophysics Report, No 113, University of Colorado.
- Kuijpers J., Pringle J., 1982, A&A 114, L7
- Korhonen T., Piirola V., Reiz A., 1984, ESO Messenger 38, 30
- Latham D.W., Liebert J., Steiner J.E., 1981, ApJ 246, 919.
- Latter W.B., Schmidt G., Green R., 1987, ApJ 320, 308
- Liller, 1977, Sky Tel. 53, 351
- Lemaitre G., Kohler D., Lacroix D. et al., 1990, A&A 228, 546
- Nauenberg M., 1972, ApJ 175, 417
- Ramsay G., Cropper M., Mason K., 1995, MN 276, 1382
- Saffer R., Wagner R., Sion E. et al., 1989, AJ 98, 668
- Schmidt G. D., Stockman H.S., Margon B., 1981, ApJ. 243, L157
- Shakhovskoy N.M., Kolesnikov S., Andronov I., 1992, In Stellar magnetism, Int. Conf. Proc., Eds Glagolevskij YU.V., Romanyuk I.I., p. 148
- Shakhovskoy N.M., Alexeev I., Andronov I., Kolesnikov S., 1993, Ann. Israel Phys. Soc. 10, 237
- Shakhovskaya, N.I., 1989, SoPH 121, 375
- Silber A., Raymond J.C., Mason P., Andronov I., Borisov N., Shakhovskoy N.M., 1996, ApJ 460, 939
- Somov, N., 1986, Astrofiz. Issled. (Izv. SAO), 22, p.73
- Somova T.A., Somov N.N., Markelov S.V., Nebelitski, 1982, In Instrumentation for astronomy with Large Optical Telescopes, Ed. C.H.Hamphries, Reidel Publ. company, p.283.
- Szkody P., Raymond J.C., Capps R.W., 1982, ApJ 257, 686
- Vikuliev N.A. et al., 1991, Astrophizich. Issled. 33, 158
- Young P., Schneider D.P., Shectman S.A., 1981, ApJ 245, 1043
- Woelk U., Beuermann K., 1992, A&A 256, 498